

РАЗДЕЛ I. ДИСФУНКЦИЯ ЭНДОТЕЛИЯ КРОВЕНОСНЫХ СОСУДОВ: ОБЩИЕ МЕДИКО-БИОЛОГИЧЕСКИЕ ВОПРОСЫ

HYPOXIC CONDITIONING SUPPRESSES CYTOTOXIC NITRIC OXIDE PRODUCTION BY ENDOTHELIUM UPON REPERFUSION FOLLOWING ACUTE MYOCARDIAL ISCHEMIA

H. Fred Downey¹, Myoung-gwi Ryou¹, Jie Sun¹, Eugenia B. Manukhina², Robert T. Mallet¹

Integrative Physiology, University of North Texas Health Science Center at Fort Worth¹, USA;

Regulatory Mechanisms of Adaptation, Institute of General Pathology and Pathophysiology², Russian Federation

There is increasing evidence that hypoxic conditioning, in particular intermittent and repetitive exposure to moderate hypoxia, induces adaptive responses that protect the heart from ischemic insults.¹ A number of mechanisms have been proposed to account for the cardioprotective responses to hypoxia stress. These mechanisms include 1) activation of ATP-sensitive K⁺ channels in cellular^{2,3} and mitochondrial⁴⁻⁶ membranes, 2) production of cardioprotective proteins including glycolytic enzymes^{7,8} and heat-shock proteins,^{9,12} 3) release of erythropoietin,¹³ 4) increased production of NO by enhanced inducible nitric oxide synthase,¹⁴⁻¹⁶ 5) angiogenesis,¹⁷ 6) attenuation of apoptosis,¹⁸ and enhancement of antioxidative defenses.¹⁹⁻²¹ Although there is evidence that hypoxia-induced increase in NO is cardioprotective,^{22,23} there is also evidence that excessive NO might exacerbate myocardial damage.^{24,25}

In the first study of hypoxia induced cardioprotection in a large animal, we found remarkable attenuation of myocardial infarct size and absence of lethal arrhythmias in hypoxia-conditioned dogs.²⁶ As shown in Figure 1, dogs that completed a program of intermittent, moderate hypoxia (9.5 -10.0% FIO₂, 5-10 min/cycle, 5-8 cycles/d with intervening 4 min normoxia for 20 d) had negligible myocardial infarction following 1 hr occlusion and 5 hr reperfusion of the left anterior descending coronary artery (LAD). Hearts of untreated control dogs had large infarctions, and the majority of these untreated animals developed ventricular tachycardia or ventricular fibrillation. Collateral blood flow to the ischemic territory did not differ among the groups, so cardioprotection by intermittent hypoxia was not due to collateral angiogenesis.




Figure 1. Intermittent, normobaric hypoxia conditioning (IHC) prevents myocardial infarction. Masses of myocardial infarct (infarct size; IS), ischemic area at risk (AAR), and left ventricular free wall + septum (LV) were measured after 1 hour occlusion of the left anterior descending coronary artery and 5 h reperfusion. Values are means \pm SEM from 7 acute non-conditioned dogs (open bars), 12 dogs hypoxia-conditioned for 20 consecutive days (IHC; cross-hatched bars), and 10 dogs sham-conditioned with normoxic air for 20 consecutive days (hatched bars). * $P < 0.001$ vs. both non-hypoxic groups.

Whereas increased myocardial NO may be a beneficial result of hypoxia conditioning, excessive release of NO by endothelium during reperfusion may be deleterious. Thus, we hypothesized that hypoxia conditioning suppresses myocardial formation of excessive NO by the endothelial NO synthase isoform (eNOS) upon coronary artery reperfusion. In the absence of hypoxic conditioning, excessive NO would react with superoxide to produce cytotoxic peroxynitrite and lead to myocardial injury.²⁷

Materials and Methods

Mongrel dogs were conditioned by a 20 d program of intermittent normobaric hypoxia as described above,²⁶ and compared with non-conditioned control dogs. One day after the conditioning program was completed, myocardium was sampled to measure left ventricular NO synthase (NOS) activity (colorimetric assay) and eNOS content (Western blot). In other anesthetized dogs, myocardial nitrite release, an index of NO formation,^{28,29} was measured in coronary venous blood during reperfusion following 1 h occlusion of the left anterior descending coronary artery (LAD).

Results

Figure 2 shows that hypoxic conditioning lowered left ventricular total NOS activity 45%, from 92 ± 8 to 43 ± 6 mU/g protein ($P < 0.01$).




Figure 2. Myocardial nitric oxide synthase (NOS) activity. Total NOS activity was measured in left ventricular myocardium from the left circumflex (LCX) and left anterior descending (LAD) coronary artery perfusion territories, and in right ventricular myocardium (RV). Values are means \pm SEM from 7 non-conditioned (control) and 6 hypoxia-conditioned (IHC) dogs. * $P < 0.05$ vs. control.

Consistent with these data is the finding that nitrite release into coronary venous blood upon reperfusion was sharply reduced in hypoxic conditioned dogs.

Figure 3 illustrates that peak nitrite release immediately after reperfusion was greatly reduced in the hypoxia conditioned dogs,

and Figure 4A shows that significantly less total nitrite was released during the first 5 min of reperfusion in these dogs. If coronary flow were reduced during reperfusion in the hypoxia-conditioned dogs, reduced endothelial shear stress might lessen NO production and decrease nitrite release. Figure 4B shows that peak flow in the reperfused LAD was similar in control and hypoxia conditioned dogs, suggesting that NO does not play an essential role in early reperfusion hyperemia; moreover, lower nitrite release in the hypoxia-conditioned dogs was not due to decreased flow-dependent shear stress. It is interesting that between 5 and 10 min of reperfusion, both flow and nitrite release were greater in the control hearts.

Figure 5 shows a Western blot for analysis of eNOS in samples of LAD perfused myocardium from three hypoxia-conditioned and four control dogs. It is apparent that eNOS content was reduced in the hypoxia-conditioned myocardium.




Figure 3. Left ventricular myocardial nitrite (NO_2^-) release. NO_2^- was measured in plasma sampled simultaneously from the aorta and interventricular coronary vein during pre-ischemic baseline (BL) and reperfusion following 60 min LAD occlusion (OCCL). Means \pm SEM from 5 experiments per group.




Figure 4. Cumulative myocardial NO_2^- release (panel A) and LAD flow (panel B) during the first 5 min of LAD reperfusion. Means \pm SEM from 5 experiments per group. * $P < 0.05$ vs. control.




Figure 5. Western blot of eNOS in left ventricular myocardium. Total protein (50 μ g) from homogenates of left ventricular myocardium was electrophoretically separated, transferred to nylon membrane, and probed with eNOS-specific antibody. Actin (Coomassie blue-stained gel) served as loading control.

Discussion

Hypoxia conditioning protects myocardium from ischemic injury. It is likely that multiple mechanisms contribute to this protection. Blocking NO synthesis can exacerbate myocardial ischemic damage,²² so NO is an important cardioprotective agent. However, moderate reductions in NO synthesis are cardioprotective,^{30,31} suggesting that beyond an optimal concentration, excessive NO may contribute to ischemic damage. This is especially likely in the setting of coronary artery reperfusion, where endothelial release of NO would be accelerated to the extent that cytotoxic peroxide and other NO derivatives might be formed in the presence of elevated superoxide.²⁷

The present findings demonstrate that hypoxic conditioning of dogs reduces myocardial NOS activity and release of nitrite into coronary venous blood at reperfusion. Since hypoxia is expected to elevate iNOS,¹⁴⁻¹⁶ the present finding of reduced total NOS activity could be explained if eNOS were concurrently reduced. Our Western blot results demonstrate that this is the case. Thus, we conclude that abruptly reperfused coronary endothelium releases excessive NO, so cytotoxic peroxynitrite and other degradation products of NO are formed. Hypoxia conditioning attenuates eNOS activity, prevents excessive NO upon reperfusion and, thus, protects myocardium from ischemia-reperfusion injury.

Acknowledgements

This work was supported by the U.S. National Institutes of Health (grants HL064785 and HL071684), the Netherlands Organization for Scientific Research (NWO grant 047-014-016), and the Russian Foundation for Basic Research (grant 03-04-49065). Arthur G. Williams, Jr. provided excellent technical support.

References

1. Manukhina EB, Downey HF, Mallet RT. Role of nitric oxide in cardiovascular adaptation to intermittent hypoxia. *Exp Biol Med*, in press.
2. Shigematsu S, Arita M. Anoxia-induced activation of ATP-sensitive K⁺ channels in guinea pig ventricular cells and its modulation by glycolysis. *Cardiovasc Res* 1997; 35:273-282
3. Liu H, Chen H, Yang X, Cheng J. ATP sensitive K⁺ channel may be involved in the protective effects of preconditioning in isolated guinea pig cardiomyocytes. *Chin Med J* 2001; 114:178-182
4. Asemu G, Papoušek F, Ošťádal B, Kolár F. Adaptation to high altitude hypoxia protects the rat heart against ischemia-induced arrhythmias. Involvement of mitochondrial K_{ATP} channel. *J Mol Cell Cardiol* 1999; 31:1821-1831
5. Neckár J, Szárszoi O, Kóten L, Papoušek F, Ošťádal B, Grover GJ, Kolár F. Effects of mitochondrial K_{ATP} modulators on cardioprotection induced by chronic high altitude hypoxia in rats. *Cardiovasc Res* 2002; 55:567-575
6. Zhu H-F, Dong J-W, Zhu W-Z, Ding H-L, Zhou Z-N. ATP-dependent potassium channels involved in the cardiac protection induced by intermittent hypoxia against ischemia/reperfusion injury. *Life Sci* 2003; 73:1275-1287
7. Bass A, Ošťádal B, Procházka J, Pelouch V, Samanek M, Stejskalova M. Intermittent high altitude-induced changes in energy metabolism in the rat myocardium and their reversibility. *Physiol Bohemoslov* 1989; 38:155-161
8. Daneshmand Z, Garcia Riera MP, Verdys M, Rossi A. Differential responses to chronic hypoxia and dietary restriction of aerobic capacity and enzyme levels in the rat myocardium. *Mol Cell Biochem* 2000; 210: 159-166
9. Deindl E, Kolár F, Neubauer E, Vogel S, Schaper W, Ošťádal B. Effect of intermittent high altitude hypoxia on gene expression in rat heart and lung. *Physiol Res* 2003; 52:147-157
10. Kacimi R, Chentoufi J, Honbo N, Long CS, Karlner JS. Hypoxia differentially regulates stress proteins in cultured myocytes: role of the p38 stress-activated kinase signaling cascade, and relation to cytoprotection. *Cardiovasc Res* 2000; 46:139-150
11. Meerson FZ, Malyshev IYu, Zamotrinsky AV. Differences in adaptive stabilization of structures in response to stress and hypoxia relate to the accumulation of hsp70 isoforms. *Mol Cell Biochem* 1992; 111:87-95
12. Zhong N, Zhang Y, Fang QZ, Zhou ZN. Intermittent hypoxia exposure-induced heat-shock protein 70 expression increases resistance of rat heart to ischemic injury. *Acta Pharmacol Sin* 2000; 21:467-472
13. Cai Z, Manalo DJ, Wei G, Rodriguez ER, Fox-Talbot K, Lu H, Zweier JL, Semenza GL. Hearts from rodents exposed to intermittent hypoxia or erythropoietin are protected against ischemia-reperfusion injury. *Circulation* 2003; 108:79-85

14. Grilli A, De Lutiis MA, Patruno A, Speranza L, Cataldi A, Centurione L, Taccardi AA, Di Napoli P, De Caterina R, Barbacane R, Conti P, Felaco M. Effect of chronic hypoxia on inducible nitric oxide synthase expression in rat myocardial tissue. *Exp Biol Med* 2003; 228:935-942
15. Jung F, Palmer LA, Zhou N, Johns NA. Hypoxic regulation of inducible nitric oxide synthase via hypoxia inducible factor-1 in cardiac myocytes. *Circ Res* 2000; 86:319-325
16. Ding H-L, Zhu H-F, Dong J-W, Zhu W-Z, Yang W-W, Yang H-T, Zhou Z-N. Inducible nitric oxide synthase contributes to intermittent hypoxia against ischemia/reperfusion injury. *Acta Pharmacol Sin* 2005; 26:315-322
17. Zhong N, Zhang Y, Zhu H-F, Wang J-C, Fang Q-Z, Zhou Z-N. Myocardial capillary angiogenesis and coronary flow in ischemia tolerance rat by adaptation to intermittent high altitude hypoxia. *Acta Pharmacol Sin* 2002; 23:305-310
18. Dong J-W, Zhu H-F, Zhu W-Z, Ding H-L, Ma T-M, Zhou Z-N. Intermittent hypoxia attenuates ischemia/reperfusion induced apoptosis in cardiac myocytes via regulating Bcl-2/Bax expression. *Cell Res* 2003; 13:385-391
19. Irlbeck M, Iwai T, Lerner T, Zimmer H-G. Effects of angiotensin II receptor blockade on hypoxia-induced right ventricular hypertrophy in rats. *J Mol Cell Cardiol* 1997; 29:2931-2939
20. Nakanishi K, Tajima F, Nakamura A, Yagura S, Ookawara T, Yamashita H, Suzuki K, Taniguchi N, Ohno H. Effects of hypobaric hypoxia on antioxidant enzymes in rats. *J Physiol* 1995; 489:869-876
21. Zhu H-F, Dong J-W, Zhu W-Z, Ding H-L, Zhou Z-N. ATP-dependent potassium channels involved in the cardiac protection induced by intermittent hypoxia against ischemia/reperfusion injury. *Life Sci* 2003; 73:1275-1287
22. Xi L, Tekin D, Gursoy E, Salloum F, Levasseur JE, Kukreja RC. Evidence that NOS2 acts as a trigger and mediator of late preconditioning induced by acute systemic hypoxia. *Am J Physiol Heart Circ Physiol* 2002; 283:H5-H12
23. Baker JE, Holman P, Kalyanaraman B, Pritchard KA Jr. Adaptation of hearts to chronic hypoxia increases tolerance to subsequent ischemia by increased nitric oxide production. *Adv Exp Med Biol* 1998; 454:203-217
24. Zhang Y, Davies LR, Martin SM, Coddington WJ, Miller FJ Jr, Buettner GR, Kerber RE. The nitric oxide donor S-nitroso-N-acetylpenicillamine (SNAP) increases free radical generation and degrades left ventricular function after myocardial ischemia-reperfusion. *Resuscitation* 2003; 59:345-352
25. Ungvari Z, Gupte SA, Recchia FA, Batkai S, Pacher P. Role of oxidative-nitrosative stress and downstream pathways in various forms of cardiomyopathy and heart failure. *Curr Vasc Pharmacol* 2005; 3:221-229
26. Zong P, Setty S, Sun W, Martinez R, Tune JD Ehrenbourg IV, Tkatchouk EN, Mallet RT, Downey HF. Intermittent hypoxic training protects canine myocardium from infarction. *Exp Biol Med* 2004; 229:806-812
27. Beckman JS, Koppenol WH. Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and the ugly. *Am J Physiol Cell Physiol* 1996; 271:C1424-C1437
28. Lauer T, Preik M, Rassaf T, Strauer BE, Deussen A, Feelisch M, Kelm M. Plasma nitrite rather than nitrate reflects regional endothelial nitric oxide synthase activity but lacks intrinsic vasodilator action. *Proc Nat Acad Sci USA* 2001; 98:12814-12819
29. Kelnibongard P, Dejam A, Lauer T, Rassaf T, Schindler A, Picker O,

Scheeren T, Godecke A, Schrader J, Schulz R, Heusch G, Schaub G, Bryan N, Feelisch M, Kelm M. Plasma nitrite reflects constitutive nitric oxide synthase activity in mammals. *Free Radic Biol Med* 2003; 35:790-796

30. Kobara M, Tatsumi T, Takeda M, Mano A, Yamanaka S, Shiraishi J, Keira N, Matoba S, Asayama J, Nakagawa M. The dual effects of nitric oxide synthase inhibitors on ischemia-reperfusion injury in rat hearts. *Basic Res Cardiol* 2003; 98:319-328

31. Yasmin W, Strynadka KD, Schulz R. Generation of peroxynitrite contributes to ischemia-reperfusion injury in isolated rat hearts. *Cardiovasc Res* 1997; 33:422-432

НО-ЗАВИСИМЫЕ МЕХАНИЗМЫ ФОРМИРОВАНИЯ КИСЛОРОДСВЯЗУЮЩИХ СВОЙСТВ ГЕМОГЛОБИНА

Зинчук В.В.

*УО «Гродненский государственный медицинский университет»,
Беларусь*

В большинстве исследований, термин "дисфункция эндотелия" используется как синоним нарушений эндотелиального метаболизма L-аргинин-NO системы, так как монооксид азота (NO) является сигнальным агентом, участвующим в опосредовании его функций и основное его количество, образуемого в организме, осуществляется в эндотелии. Снижение синтеза NO может быть вызвано: отсутствием исходного субстрата - L-аргинина, отсутствием кофакторов, сниженной экспрессией фермента и повышением уровня эндогенных ингибиторов фермента (например, асимметрично диметилированных производных аргинина). Кроме того, снижение антиоксидантной способности организма также может вести к зависимой от NO дисфункции эндотелия, так как снижение его эффектов может вызываться усиленной его инактивацией в результате повышенной выработки кислородных радикалов.

В последние годы проблема изучения физиологических эффектов NO приобрела новый аспект, а именно его взаимодействие с различными компонентами крови, и в частности, с гемоглобином. Предложена теория сосудистого «сохранения» и транспорта NO гемоглобином. В литературе широко обсуждались различные стороны этого вопроса, что получило отражение в ряде публикаций с весьма интригующими заголовками: «Гемоглобин: транспортер NO, инактиватор NO или ни то ни другое?» [Hobbs A.J. et al., 2002]; «Реакции азота с гемоглобином: взгляд через шторм SNO» [Gladwin M.T. et al., 2003];